












#### Water from Snow

About 75% of the Western US freshwater supply comes from snowmelt.

Reservoirs in California hold about a year and a half of annual runoff, making for strong sensitivity to interannual variability in snowfall

The definition of optimization of reservoir storage, hydroelectric generation, achievement of environmental metrics depends on the timing and magnitude of runoff and varies from basin to basin


Snowmelt modeling and forecasting is migrating to physically based models, ultimately demanding markedly better snow information



# This is how we have known SNOW WATER EQUIVALENT



#### Operational snow monitoring









#### Snow Density Modeling

Mean error -26 kg/m<sup>3</sup>

MAE 31 kg/m<sup>3</sup>


Elevation bias







SWE very low is when densities fall apart -









# ASO-DM1 Results





# ASO-DM1 Results



# ASO-DM1 Results









# Hetch Hetchy Operations


balancing streamflow, water supply, & hydropower needs








#### Tuolumne Basin above Hetch Hetchy Res. SWE/Met Stations & PRMS Model Units




aggregated to PRMS model units



### Hydro modeling results


Hetch Hetchy Reservoir Tuolumne River, CA



- Precipitation and temperature were forecasted from 6/1
- Simulated and observed inflow are similar from 5/15 through 6/1, the forecast date
- Modeled SWE (red) corrected by ASO SWE (blue)

# Hydro modeling results


Hetch Hetchy Reservoir Tuolumne River, CA



- Observed inflow 28 TAF less than initial prediction
- ASO-corrected forecast closer to observed
- Reservoir operations adjusted (draft reduced) after ASO forecast
  - HH reached top of gates, was full
  - no water lost to spill

# SWE/Runoff comparison

- Total Snow Water declined from 218 to 15 TAF in 65 days
- Reservoir storage reached capacity as last snow melted
- Melt rates peaked on May 1 based on solar input and snow extent, and then declined as snow extent decreased



#### ASO in 2014+

- ♦ We begin flying the Tuolumne on ~23 March
- Quantification of Tuolumne SWE should be available 24 March, weather permitting
- This spring is likely to be endmember snow season
- Rim Fire of 2013 in and near Yosemite is a science target for California, ecosystem scientists, and hydrologists. Our flights are key to understanding hydrologic response
- Implementation of new density modeling and physical constraints on density modeling
- Scaling to entire Sierra Nevada in coming years.